

What is claimed is:

1. A method for augmenting an immune response in a patient comprising the step of administering an amount of flt3-ligand to the patient sufficient to generate an increase in the number of the patient's dendritic cells.
2. A method according to claim 1, further comprising the step of administering one or more of the molecules selected from the group consisting of GM-CSF, IL-4, TNF- α , IL-3, c-kit ligand, and fusions of GM-CSF and IL-3.
3. A method for augmenting an immune response in a patient having an infectious disease, comprising the step of administering of administering flt3-ligand in an amount sufficient to generate an increase in the number of the patient's dendritic cells.
4. A method according to claim 3, further comprising the step of administering one or more of the molecules selected from the group consisting of GM-CSF, IL-4, TNF- α , IL-3, c-kit ligand, and fusions of GM-CSF and IL-3.
5. A method according to claim 3, wherein the infectious disease is HIV.
6. A method for augmenting an immune response in a patient having a cancerous or neoplastic disease, comprising the step of administering flt3-ligand in an amount sufficient to generate an increase in the number of the patient's dendritic cells.
7. A method according to claim 6, further comprising the step of administering one or more of the molecules selected from the group consisting of GM-CSF, IL-4, TNF- α , IL-3, c-kit ligand, and fusions of GM-CSF and IL-3.
8. A preparation of dendritic cells having at least two cell surface markers selected from the group consisting of CD1a, HLA-DR and CD86, produced by contacting hematopoietic stem or progenitor cells with flt3-ligand.
9. A dendritic cell preparation according to claim 8 produced further by contacting the hematopoietic stem or progenitor cells with a molecule selected from the group consisting of GM-CSF, IL-4, TNF- α , IL-3, c-kit ligand, and fusions of GM-CSF and IL-3.

10. An antigen-expressing dendritic cell population produced by the process of
- (a) contacting hematopoietic stem or progenitor cells with flt3-ligand in an amount sufficient to generate a dendritic cell population;
 - (b) either (i) exposing the dendritic cells to an antigen-specific peptide or (ii) transfecting the dendritic cells with a gene encoding an antigen-specific peptide;
 - (c) allowing the dendritic cells to process and express the antigen; and
 - (d) purifying the antigen-expressing dendritic cells.
11. A dendritic cell population according to claim 10 wherein step (a) of the process further comprises contacting the hematopoietic stem or progenitor cells with a molecule selected from the group consisting of GM-CSF, IL-4, TNF- α , IL-3, c-kit ligand, and fusions of GM-CSF and IL-3.
12. A method of driving hematopoietic stem or progenitor cells to a dendritic cell lineage comprising contacting such hematopoietic stem or progenitor cells with flt3-ligand.
13. A method of preparing an antigen-presenting dendritic cell population comprising the steps of:
- (a) contacting hematopoietic stem or progenitor cells with flt3-ligand in an amount sufficient to generate a dendritic cell population;
 - (b) either (i) exposing the dendritic cells to an antigen-specific peptide or (ii) transfecting the dendritic cells with a gene encoding an antigen-specific peptide;
 - (c) allowing the dendritic cells to process and express the antigen; and
 - (d) purifying the antigen-expressing dendritic cells.
14. A method according to claim 13, wherein step (a) further comprises contacting the hematopoietic stem or progenitor cells with a molecule selected from the group consisting of GM-CSF, IL-4, TNF- α , IL-3, c-kit ligand, and fusions of GM-CSF and IL-3.
15. A method of preparing antigen-specific T cells comprising the steps of:
- (a) contacting hematopoietic stem or progenitor cells with flt3-ligand in an amount sufficient to generate a dendritic cell population;

- (b) either (i) exposing the dendritic cells to an antigen-specific peptide or (ii) transfecting the dendritic cells with a gene encoding an antigen-specific peptide;
- (c) allowing the dendritic cells to process and express the antigen; and
- (d) allowing the dendritic cells to present the antigen to T cells.

16. A method of enhancing a mammal's immune response to a vaccine antigen, comprising the steps of administering to such mammal an immunogenic amount of the vaccine antigen and an immunogenicity-augmenting amount of flt3-ligand in concurrent or sequential combination with such vaccine antigen.

17. A vaccine adjuvant comprising a molecule selected from the group consisting of c-kit ligand and flt3-ligand.

18. A method for inducing tolerance of graft tissue in a host, comprising administering flt3-ligand to the host in an amount sufficient to increase the number of dendritic cells.

19. A dendritic cell expansion media comprising an effective amount of flt3-ligand and a cytokine selected from the group consisting of IL-3, IL-4, GM-CSF, TNF, C-Kit ligand and GM-CSF/IL-3 fusion proteins.